



# 2022 Kenya Demographic and Health Survey Fact Sheet

## Kisii County

Characteristics of Households and Respondents	Kisii	Kenya
Household population with access to at least basic drinking water service (%)	44	68
Household population with at least basic sanitation service (%)	29	41
Household population relying on clean fuels and technologies for cooking, space heating, & lighting (%)	10	21
Women age 15-49 with no formal education (%)	1	6
Men age 15-49 with no formal education (%)	<1	3
<b>Fertility and Family Planning (FP)</b>		
Total fertility rate (number of children per woman)	3.0	3.4
Teenage pregnancy (% age 15-19 who have ever been pregnant)	14	15
Use of modern method of FP (% of married women age 15-49)	64	57
Unmet need for FP <sup>1</sup> (% of married women age 15-49)	15	14
Demand for FP satisfied by modern methods (% of married women age 15-49)	76	75
<b>Maternal and Child Health</b>		
Births delivered by a skilled provider <sup>2</sup> (%)	94	89
Women age 15-49 who had a live birth and had 4+ antenatal visits (%)	63	66
Women age 15-49 with a postnatal check during the first 2 days after birth (%)	89	78
Births with a postnatal check during the first 2 days after birth (%)	91	83
Children age 12-23 months fully vaccinated (basic antigens) <sup>3</sup> (%)	90	80
Neonatal mortality <sup>4</sup> (deaths per 1,000 live births)	21	21
Infant mortality <sup>4</sup> (deaths per 1,000 live births)	24	32
Under-5 mortality <sup>4</sup> (deaths per 1,000 live births)	40	41
<b>Nutrition</b>		
Children under 5 who are stunted (%) (too short for their age)	16	18
Children under 5 who are wasted (%) (too thin for their height)	3	5
Children under 5 who are underweight (%) (too thin for their age)	8	10
Children under 5 who are overweight (%) (too heavy for their age)	6	3
<b>Malaria</b>		
Household population with access to an insecticide-treated net (ITN) (%)	79	50
Household population who slept under an ITN the night before the survey (%)	71	43
Women age 15-49 who received 3 or more doses of SP/Fansidar during pregnancy (%)	11	13 <sup>a</sup>
<b>Gender-based Violence</b>		
Women age 15-49 who have ever experienced physical violence since age 15 (%)	36	34
Women age 15-49 who experienced physical violence in the last 12 months (often or sometimes) (%)	20	16
Women age 15-49 who have ever experienced sexual violence (%)	16	13
Women age 15-49 who experienced sexual violence in the last 12 months (%)	10	7
Women age 15-49 who have ever been married or had an intimate partner & have ever experienced physical, sexual, or psychological/emotional violence committed by their most recent husband/partner (%)	53	40
Women age 15-49 who have ever been married or had an intimate partner & have experienced physical, sexual, or psychological/emotional violence committed by any husband/partner in the last 12 months (%)	41	28

Footnotes: <sup>1</sup> Women who do not want any more children or want to wait at least 2 years before their next birth but are not currently using a method of FP. <sup>2</sup> Skilled provider is a doctor, nurse, midwife, or clinical officer. <sup>3</sup> Basic antigens include BCG, rotavirus, DPT-HepB-Hib, pneumococcal, OPV, IPV, and MR. <sup>4</sup> County-level childhood mortality rate estimates are for the 10-year period before the survey and national level childhood mortality rate estimates are for the 5-year period before the survey. <sup>a</sup> Kenya total is among counties where IPTp is not implemented. Kisii County does not implement IPTp.

